

## RESEARCH ARTICLE

## ***In vitro* Evaluation of Cytotoxic Activities of Essential Oil from *Moringa oleifera* Seeds on HeLa, HepG2, MCF-7, CACO-2 and L929 Cell Lines**

Elsayed Ahmed Elsayed<sup>1,2\*</sup>, Mahmoud A. Sharaf-Eldin<sup>3,4</sup>, Mohammad Wadaan<sup>1</sup>

### **Abstract**

*Moringa oleifera* Lam. (Moringaceae) is widely consumed in tropical and subtropical regions for their valuable nutritional and medicinal characteristics. Recently, extensive research has been conducted on leaf extracts of *M. oleifera* to evaluate their potential cytotoxic effects. However, with the exception of antimicrobial and antioxidant activities, little information is present on the cytotoxic activity of the essential oil obtained from *M. oleifera* seeds. Therefore, the present investigation was designed to investigate the potential cytotoxic activity of seed essential oil obtained from *M. oleifera* on HeLa, HepG2, MCF-7, CACO-2 and L929 cell lines. The different cell lines were subjected to increasing oil concentrations ranging from 0.15 to 1 mg/mL for 24h, and the cytotoxicity was assessed using MTT assay. All treated cell lines showed a significant reduction in cell viability in response to the increasing oil concentration. Moreover, the reduction depended on the cell line as well as the oil concentration applied. Additionally, HeLa cells were the most affected cells followed by HepG2, MCF-7, L929 and CACO-2, where the percentages of cell toxicity recorded were 76.1, 65.1, 59.5, 57.0 and 49.7%, respectively. Furthermore, the IC<sub>50</sub> values obtained for MCF-7, HeLa and HepG2 cells were 226.1, 422.8 and 751.9 µg/mL, respectively. Conclusively, the present investigation provides preliminary results which suggest that seed essential oil from *M. oleifera* has potent cytotoxic activities against cancer cell lines.

**Keywords:** Anticancer agents - cytotoxicity - medicinal plants - *Moringa* - seed oil

*Asian Pac J Cancer Prev*, 16 (11), 4671-4675

### **Introduction**

Cancer is one of the most harmful diseases leading to human death in developing as well as developed countries. In 2012, 14.1 million cancer cases have been reported worldwide, from which 52.5% cases occurred in men and 45.4% cases were in women (Sultana et al., 2014). The number of cancer cases is estimated to increase by about 70% in 2035 (Ferlay, 2014). Breast and cervical cancers are two of the most common cancer causes in women worldwide (WCR, 2008; Farshori et al., 2013; Srisuwan et al., 2014). In Saudi Arabia, there was a significant increase in the age-standardized incidence rate (ASIR) female breast cancer between 2001 and 2008, with the eastern province recording the highest overall ASIR, at 26.6 per 100,000 women (Alghamdi et al., 2013). On the other hand, the occurrence of cervical cancer in Saudi Arabia is very low, ranking 11<sup>th</sup> between all female cancers, and accounts only for 2.4% of all new cases reported (Alsbeih et al., 2013). Colon cancer, the third most common cancer worldwide, accounts for the fourth occurring cancer among all age groups in Saudi Arabia (Al-Kuraya et al.,

2006). Liver cancer is considered as the third cause of cancer mortality worldwide (Farooq et al., 2013). In Saudi Arabia, liver cancer accounts for about 10 and 4% of total cancers in males and females, respectively (Al-Ahmadi and Al-Zahrani, 2013).

Standard cancer treatment protocols depended largely on chemotherapy and/or radiotherapy, as well as surgical intervention (Huang et al., 2014). However, due to sever physical and psychological side effects of such standard treatment methods as well as recent advancements and developments in the field of isolation and identification of phytochemical compounds, there have been an increasing trend during the last two decades toward the application of traditional herbal medicine as a potential source for anticancer agents (Awodele et al., 2012; Shahat et al., 2013). Nowadays, several anticancer agents, currently used in clinical trials, have been isolated and identified from plants and their constituents. Vinblastine, camptothecin, homoharringtonine, ellipticine and paclitaxel have been isolated from *Catharanthus roseus* G. Don. (Apocynaceae), *Camptotheca acuminata* Decne (Nyssaceae), *Cephalotaxus harringtonia* var.

<sup>1</sup> Department of Zoology, Bioproducts Research Chair, Faculty of Science, King Saud University, Riyadh, <sup>3</sup>Sara bint Rached bin Ghonaim Research Chair for Cultivating Non-traditional Medicinal and Aromatic Plants, Biology Department, College of Science and Humanities, Prince Sattam bin Abdulaziz University, Alkharj, Kingdom of Saudi Arabia, <sup>2</sup>Department of Natural and Microbial Products, National Research Centre, Dokki, <sup>4</sup>Department of Medicinal and Aromatic Plants Research, National Research Centre, Cairo, Egypt \*For correspondence: eaelsayed@ksu.edu.sa

*drupacea* (Sieb and Zucc.) (Cephalotaxaceae), *Bleekeria vitensis* A.C. Sm. and *Taxus brevifolia* Nutt. (Taxaceae), respectively (Bakar et al., 2010; Kuete et al., 2011; Sultana et al., 2014).

The flowering plant family Moringaceae contains only the genus *Moringa* with 13 species distributed from tropical to subtropical regions and ranging in size from tiny herbs to massive trees (Janick and Paull, 2008). The most widely known species is the *M. oleifera* Lam., also known as horseradish tree, drumstick tree, melonge or malunggay (Awodele et al., 2012). Beside their valued nutritional characteristics, different parts of *M. oleifera* (leaves, flowers, park, pods, stem, roots and seeds) have been reported to possess different biological properties, i.e. antimicrobial, antitrypanosomal, hypotensive, antiulcer, hypocholesterolemic, antispasmodic, antioxidant, anti-inflammatory as well as anti-cancer (Sreelatha et al., 2011; Awodele et al., 2012; Satish et al., 2013; Tiloke et al., 2013; Vongsak et al., 2013a; 2013b; Hannan et al., 2014). *M. oleifera* seeds contain 33-41% w/w vegetable oil (Rashid et al., 2008). The seed oil of *M. oleifera* resembles olive oil and is widely used in cooking, lighting, hairdressing as well as soap and perfume industries (Ghazali and Mohammed, 2011). Moreover, it has been found to exhibit antimicrobial and antioxidant activities (Siddhuraju and Becker, 2003; Chuang et al., 2007).

Taking into consideration the lack of anticancer studies on seed oils from *Moringa*, to the best of our knowledge, the present investigation was designed to evaluate the possible anti-proliferative activity of seed oil obtained from *M. oleifera* on human cervical, hepatocellular, breast, colon cancers as well as mouse fibroblast cell lines.

## Materials and Methods

### Chemicals and reagents

Dulbecco's Modified Eagle's Medium (DMEM) culture medium containing 4.5 g/L glucose was purchased from AppliChem, Darmstadt, Germany. Fetal bovine serum (FBS), 0.25% Trypsin-EDTA solution and antibiotic/antimycotic solution were purchased from GIBCO® Invitrogen, Life Technologies, USA. Other chemicals and reagents were of cell culture grade and were purchased from Sigma-Aldrich Chemical Company, St. Louis, MO, USA. Disposable culture ware and consumable materials were procured from Corning, NY, USA.

### Preparation of seed oils

Commercial purified oil of *M. oleifera* seeds was kindly provided by AsiaNet Trading Est., Riyadh, KSA. The oils were extracted by cold pressing in order to retain all the aromatic, chemical and nutritional characteristics of the oil. The purified oil was used to prepare a stock solution of 1 mg/mL in DMSO. The working seed oil solutions were prepared by serially diluting the stock oil solution with the cultivation medium. Finally, the working solution sets were filtered using 0.22 µm sterile syringe filters (Millipore, USA).

### Cell lines and cultivation conditions

HeLa, human cervical cancer; HepG2, human

hepatocellular carcinoma; MCF-7, human breast cancer; CACO-2, Caucasian colon adenocarcinoma and L929, mouse fibroblast cell lines, were purchased from Sigma-Aldrich Chemical Company, USA. Cells were cultivated in DMEM, supplemented with 10% FBS, 100x, 1% antibiotic/antimycotic solution and 3.6 g/L NaHCO<sub>3</sub>. Cells were routinely sub-cultured and propagated in a humidified CO<sub>2</sub> incubator (ShelLab, USA) at 5% CO<sub>2</sub>, 37°C and 95% humidity. Upon testing the effect of seed oil on cell viability, cells were trypsinized, centrifuged and washed twice with sterile PBS buffer solution. The viable cell count and viability were estimated using Trypan blue exclusion method (Al-Sheddi et al., 2014). Cell batches with viabilities above 95% were used to run the cytotoxicity assays.

### Treatments and experimental design

The tested different cell lines were used to evaluate the cytotoxic effects of *M. oleifera* seed oil. Cells were treated with different concentrations of the prepared seed oil ranging from 15.6 µg/mL to 1 mg/mL. After cell treatment, the cytotoxic activities of the tested seed oils were screened using 3-(4,5-dimethylthiazol-2-yl),5-biphenyl tetrazolium bromide (MTT). Control sets were treated with DMSO only (final concentration not exceeding 0.5%). In addition, the morphological changes were assessed using phase contrast inverted microscope.

### Cytotoxicity assay

The percentage of cell viability was evaluated with MTT according to the method described by Siddiqui et al. (2008). In brief, after trypsinization and washing, cells were resuspended and were used to inoculate 96 well culture plates at a final concentration of 10<sup>4</sup> cells/100 µl/well. The plates were incubated at 5% CO<sub>2</sub> and 37°C to allow cell adherence. After 24h, the medium was replaced with fresh medium containing different concentrations from the prepared seed oil serial dilutions, and were allowed to grow for another 24h. After incubation, 10 µl MTT (5 mg/mL in PBS) were added to each well, and the plates were further incubated for 4h. The supernatants were then discarded and 200 µl of DMSO were added to each well and mixed to dissolve the precipitated dark blue formazan crystals. The absorbance of formazan, directly proportional to the number of living cells, was read at 550 nm using a microplate reader (Thermo Scientific, USA). The percentages of viability (%V) and inhibition (%I) were calculated as:

$$\%V=100(A_t/A_c)$$

$$\%I=100 [1-(A_t/A_c)]$$

where  $A_t$  and  $A_c$  are the absorbances for treated and control cells, respectively. IC<sub>50</sub> values were defined as sample concentration inhibiting 50% of cell growth, and they were obtained from the linear regression of the calibration curve.

### Morphological examination

Cells treated with different concentrations of seed oil were microscopically examined for morphological changes using an inverted contrast microscope (Nikon Eclipse T500, Japan) at 10x magnification.

### Statistical analysis

The obtained data were analyzed with the help of SPSS 9.0, and the results were given as mean±SD of three replicates. The mean comparison between different evaluated groups was performed using ANOVA one-way analysis of variance. Statistical significance was defined when  $p < 0.05$ .

## Results

### Cytotoxicity assay

The cytotoxic effect of the *M. oleifera* essential oil on different cell lines (HeLa, HepG2, MCF-7, CACO-2 and L929) has been evaluated using standard MTT assay. The different cell lines have been treated with increasing concentrations (15-1000  $\mu\text{g/mL}$ ) for 24h. The obtained results illustrated in Figure 1A-E show clearly, that increasing the concentration of the essential oil adversely decreased the viability of the treated cell lines, with a highly significant effect ( $p < 0.001$ ). Moreover, the highest concentration applied (1 mg/mL) gave the highest decrease in cell viability in all cell lines, where the viability recorded  $23.9 \pm 0.85$ ,  $34.93 \pm 0.32$ ,  $40.48 \pm 6.61$ ,  $50.28 \pm 5.86$  and  $42.99 \pm 0.17$  for HeLa, HepG2, MCF-7, CACO-2 and L929 cells, respectively. Furthermore, the obtained results, for all cell lines, also revealed that the effect on cell viability was not statistically significantly changed upon treating the cells with either 15.6 and 31.3  $\mu\text{g/mL}$  or 62.5 and 125  $\mu\text{g/mL}$ . Additionally, it can be clearly observed that treatment of HeLa and MCF-7 cells with either 31.3 or 62.5  $\mu\text{g/mL}$  showed no statistically significant difference. On the other hand, CACO-2 cells showed no statistically significant difference in cell viability when treated with a concentration ranging from 31.3 to 250  $\mu\text{g/mL}$ .

The results in Figure 1F compares between the highest percentages of toxicity obtained for different cell lines at 1 mg/mL of essential oil. It can be clearly noted that HeLa cells were the most affected cells followed by HepG2, MCF-7, L929 and CACO-2, where the percentages of cell toxicity recorded 76.05, 65.07, 59.52, 57.02 and 49.72%, respectively.


The concentration of essential oil required to inhibit 50% of the cells is recorded in Table 1. The results showed that the  $\text{IC}_{50}$  value obtained for MCF-7 cells was 226.1  $\mu\text{g/mL}$ , while the values recorded for HeLa and HepG2 cells were higher by 1.9- and 3.3-folds. On the other hand, the  $\text{IC}_{50}$  values for CACO-2 and L929 cells were higher than 1 mg/mL.

### Morphological examination


Figure 2 shows the morphological changes occurred in HepG2, MCF-7, L929 as well as CACO-2 cells upon treatment with different concentrations of *M. oleifera* seed oil. Increasing the seed oil concentration ( $>125 \mu\text{g/mL}$ ) resulted in a drastic change in the morphological characteristics of the tested cell lines, which was proportional to the applied concentration. Cells started to shrink and lose their capacity to adhere to the surface of the cultivation plate. Moreover, at the highest applied concentration, cells appeared rounded and were completely floated in comparison to the control morphology.

**Table 1.  $\text{IC}_{50}$  Values Obtained for Different Cell Lines as Affected by Essential Oil From *M. oleifera* Seeds.**

L929	CACO-2	MCF-7	HepG2	HeLa	Cell Line
$>1000.0$	$>1000.0$	226.1	751.9	442.8	$\text{IC}_{50}$ ( $\mu\text{g/mL}$ )



**Figure 1. Effect of Different Concentrations of *M. oleifera* Seed Oil on the Viability of Different Cell Lines (A: HeLa, B: HepG2, C: MCF-7, D: CACO-2, E: L929, F: Toxicity on Different Cell Lines at 1 mg/mL).** Data are expressed as means±SD. Dissimilar alphabets within the same cell line indicate significantly different at  $p < 0.05$  using one-way ANOVA



**Figure 2. Morphological Changes in Different Cell Lines as Affected by Different Concentrations of *M. oleifera* Seed Oil after 24h.** Images are captured using inverted contrast microscope (Nikon Eclipse T500) at 10x magnification

## Discussion

Natural medicinal plants and their constituents provide the pharmaceutical industry with more than 60% of the compounds exhibiting potent cytotoxic activities against cancer cells (Cragg and Newman, 2005). *M. oleifera*, growing in tropical and subtropical climates, has been highly appreciated for its nutritional benefits. The leaves of *M. oleifera* are rich in proteins, minerals and vitamins A, B and C, and they contain all essential amino acids (Janick



and Paull, 2008). During the past two decades, *M. oleifera* leaves have been shown to possess different biological activities; antimicrobial, anti-inflammatory, antioxidant, anti-parasitic as well as anticancer (Gasparotto et al., 2001; Anwar et al., 2007; Ferreira et al., 2011; Awodele et al., 2012). Although *M. oleifera* has been widely investigated for its anticancer activity, scarce information can be found in the literature regarding the cytotoxic activity of its seed essential oil. Therefore, the present investigation has been carried out to evaluate the anti-proliferative activity of *M. oleifera* seed oil on different cell lines; i.e. HeLa, HepG2, MCF-7, CACO-2 and L929. Our results showed that increasing the concentration of seed oil greatly decreased the viability of different cell lines depending on the concentration applied after 24h of incubation. These results are in accordance with those of Al-Oqail et al. (2013), who obtained a dose-dependent response of different concentrations of fenugreek seed oil on HEP2, MCF-7, WISH and Vero cells. Moreover, our obtained results showed variation among different cell lines when treated with different concentrations of seed oil. In such case, HeLa cells showed the highest decrease in cell viability followed by HepG2, MCF-7, L929 and CACO-2. The obtained results are in agreement with those of Heo et al. (2014), who also evaluated anticancer effects of different parts of indigo plant on different cell lines (Hek-293, HCT-116, HeLa, MCF-7, Hep3B, SNU-1066 and SNU-601). They concluded that the plant extract effectively reduced the cell viability of different cell lines depending on the extract concentration as well as cell type. Generally, the cytotoxic effects of the essential oil obtained from *M. oleifera* seeds can be attributed to the presence, nature and composition of the oil. Saturated and unsaturated fatty acids constitute 49.1 and 50.9% of *M. oleifera* seed essential oil (Abdulkarim et al., 2005). Among these fatty acids, octadecenoic, hexadecanoic, octadecanoic and docosanoic acids correspond for 70.0, 6.8, 6.5 and 5.8%, respectively. Recently, unsaturated fatty acids have been reported to exhibit anticancer activity against many malignant cell lines *in vitro*, as well as reducing the tumor growth and cancer incidence in animal models (Schein, 2009). Furthermore, Mustafa et al. (2004) investigated the anticancer activities of different fatty acid analogs of podophyllotoxin on SK-MEL, KB, BT-549, SK-OV-3 and HL-60 cell lines. Their fatty acid analogs incorporated 10-hydroxydecanoic, 12-hydroxydodecanoic, 15-hydroxypentadecanoic, 16-hydroxyhexadecanoic, 12-hydroxyoctadec-Z-9-enoic, eicosa-Z-5,8,11,14-tetraenoic, eicosa-Z-8,11,14-trienoic, eicosa-Z-11,14-dienoic, eicosa-Z-11-enoic and eicosanoic acids. They found that their analogs, except for eicosa-Z-11-enoic and eicosanoic acids, exhibited cytotoxic effects against the different cancer cell lines. Additionally, Ravichandran and Johnson (1999) suggested that the antitumor effect of essential fatty acids is related to the increased lipid peroxidation leading to increased production of free radicals, which finally lead to the modification of membrane fatty acid composition of tumor cells.

The effect of different concentrations of seed oil on the morphology of different cell lines showed a concentration

dependent-effect where the cells started to lose their normal morphological characteristics as well as adhering capacity, and finally leading to cell detachment and death. These results are in accordance with those of Vijayarathna and Sreenivasan (2012), who evaluated the cytotoxic effects of *Elaeis guineensis* on MCF-7 and Vero cells. They found that both cell lines showed morphological alterations with cell blebbing and vacuolation and suggested autophagy as a mechanism for cell death.

Finally, from the aforementioned results, it can be concluded that seed essential oil of *M. oleifera* has statistically significant cytotoxic effects on HeLa, HepG2, MCF-7, CACO-2 and L929 cells. Moreover, the effect was proportional to the applied concentration, with HeLa, HepG2 and MCF-7 cell lines as the most affected cell lines. Additionally, the present work provides a preliminary platform for further investigation of the possible mechanism and role of *M. oleifera* seed oil on cancer cell lines.

## Acknowledgements

The Authors are thankful for the Deanship of Scientific Research at Prince Sattam bin Abdulaziz University, for supporting this work through the research project grant number 2014-01-2433.

## References

- Abdulkarim SM, Lai OM, Muhammad SKS, Long K, Ghazali HM (2005). Some physico-chemical properties of *Moringa oleifera* seed oil extracted using solvent and aqueous enzymatic methods. *Food Chem*, **93**, 253-63.
- Al-Ahmadi K, Al-Zahrani A (2013). Spatial autocorrelation of cancer incidence in Saudi Arabia. *Int J Environ Res Public Health*, **10**, 7207-28.
- Alghamdi IG, Hussain II, Alghamdi MS, El-Sheemy MA (2013). The incidence rate of female breast cancer in Saudi Arabia: an observational descriptive epidemiological analysis of data from Saudi cancer registry 2001-2008. *Breast Cancer: Targets Therapy*, **5**, 103-9.
- Al-Kuraya KS, Bavi PP, Ezzat AA, et al (2006). Colorectal carcinoma from Saudi Arabia: Analysis of MLH-1, MSH-2 and p53 genes by immunohistochemistry and tissue microarray analysis. *Saudi Med J*, **27**, 323-8.
- Al-Oqail MM, Farshori NN, Al-Sheddi ES, et al (2013). *In vitro* cytotoxic activity of seed oil of fenugreek against various cancer cell lines. *Asian Pac J Cancer Prev*, **14**, 1829-32.
- Alsbeih G, Al-Harbi N, El-Sebaie M, Al-Badawi I (2013). HPV prevalence and genetic predisposition to cervical cancer in Saudi Arabia. *Infect Agents Cancer*, **8**, 15.
- Al-Sheddi ES, Farshori NN, Al-Oqail MM, et al (2014). Cytotoxicity of *Nigella Sativa* seed oil and extract against human lung cancer cell line. *Asian Pac J Cancer Prev*, **15**, 983-7.
- Anwar F, Latif S, Ashraf M, Gilani AH (2007). *Moringa oleifera*: a food plant with multiple-medicinal uses. *Phytother Res*, **21**, 17-25.
- Awodele O, Oreagbaa IA, Odomaa S, Teixeira da Silva JA, Osunkaluc VO (2012). Toxicological evaluation of the aqueous leaf extract of *Moringa oleifera* Lam. (*Moringaceae*). *J Ethnopharmacol*, **139**, 330-6.
- Bakar MFA, Mohamad M, Rahmat A, Burr SA, Fry JR (2010). Cytotoxicity, cell cycle arrest, and apoptosis in breast

- cancer cell lines exposed to an extract of the seed kernel of *Mangifera pajang* (bambangan). *Food Chem Toxicol*, **48**, 1688-97.
- Chuang P-H, Lee C-W, Chou J-C, et al (2007). Anti-fungal activity of crude extracts and essential oil of *Moringa oleifera* Lam. *Bioresour Technol*, **98**, 232-6.
- Cragg GM, Newman DJ (2005). Plant as source of anticancer agents. *J Ethnopharmacol*, **100**, 72-9.
- Farooq M, Hozzein WN, Elsayed EA, et al (2013). Identification of histone deacetylase I protein complexes in liver cancer cells. *Asian Pac J Cancer Prev*, **14**, 915-21.
- Farshori NN, Al-Sheddi ES, Al-Oqail MM, et al (2013). Anticancer activity of *Petroselinum sativum* seed extracts on MCF-7 human breast cancer cells. *Asian Pac J Cancer Prev*, **14**, 5719-23.
- Ferlay J, Soerjomataram I, Dikshit R, et al (2014). Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Inter J Cancer*, **136**, 359-86.
- Ferreira RS, Napoleao TH, Santos AF, et al (2011). Coagulant and antibacterial activities of the water-soluble seed lectin from *Moringa oleifera*. *Lett Appl Microbiol*, **53**, 186-92.
- Gasparotto JrA, Gasparotto FM, Lourenco ELB, et al (2011). Anti-hypertensive effects of isoquercetin and extracts from *Tropaeolum majus* L: evidence for the inhibition of angiotensin converting enzyme. *J Ethnopharmacol*, **134**, 363-72.
- Ghazali HM, Mohammed AS (2011). *Moringa (Moringa oleifera)* seed oil: composition, nutritional aspects and health attributes. in 'nuts & seeds in health and disease prevention', Eds. Preedy V, Watson R and Patel V. Elsevier Academic Press, USA pp 787-93.
- Hannan Md.A, Kang J-Y, Mohibullah Md, et al (2014). *Moringa oleifera* with promising neuronal survival and neurite outgrowth promoting potentials. *J Ethnopharmacol*, **152**, 142-50.
- Heo B-G, Park Y-J, Park Y-S, et al (2014). Anticancer and antioxidant effects of extracts from different parts of indigo plant. *Indust Crops Products*, **56**, 9-16.
- Huang Y-J, Zhang Y-Y, Liu G, et al (2014). Preliminary evaluation of the *in vitro* efficacy of 1, 2-di(quinazolin-4-yl) diselane against SiHa cervical cancer cells. *Asian Pac J Cancer Prev*, **15**, 6301-6.
- Janick J, Paull RE (2008). The encyclopedia of fruit & nuts. CABI, Cambridge, UK pp 509-10.
- Kuete V, Krusche B, Youns M, et al (2011). Cytotoxicity of some Cameroonian spices and selected medicinal plant extracts. *J Ethnopharmacol*, **134**, 803-12.
- Mustafa J, Khan SI, Ma G, Walker LS, Khana IA (2004). Synthesis and anticancer activities of fatty acid analogs of podophyllotoxin. *Lipids*, **39**, 167-72.
- Rashid U, Anwar F, Moser BR, Knothe G (2008). *Moringa oleifera*: A possible source for biodiesel. *Bioresour Technol*, **99**, 8175-9.
- Ravichandran D, Johnson CD (1999). Anticancer effects of essential fatty acids. in 'pancreatic disease', Eds. Johnson CD. Springer-Verlag, London pp 325-37.
- Satish A, Kumar RP, Rakshith D, Satish S, Ahmed F (2013). Antimutagenic and antioxidant activity of *Ficus benghalensis* stem bark and *Moringa oleifera* root extract. *Inter J Chem Anal Sci*, **4**, 45-8.
- Schein DE (2009). Cytotoxicity of unsaturated fatty acids in fresh human tumor explants: concentration thresholds and implications for clinical efficacy. *Lipids Health Dis*, **8**, 54.
- Shahat AA, Alsaid MS, Alyahya MA, Higgins M, Dinkova-Kostova AT (2013). NAD(P)H: Quinone oxidoreductase 1 inducer activity of some Saudi Arabian medicinal plants. *Planta Med*, **79**, 459-64.
- Siddhuraju P, Becker K (2003). Antioxidant properties of various solvent extracts of total phenolic constituents from three different agro-climatic origins of drumstick tree (*M. oleifera* Lam.). *J Agric Food Chem*, **51**, 2144-55.
- Siddiqui MA, Singh G, Kashyap MP, et al (2008). Influence of cytotoxic doses of 4-hydroxynonenal on selected neurotransmitter receptors in PC-12 cells. *Toxicol in vitro*, **22**, 1681-8.
- Sreelatha S, Jeyachitra A, Padma PR (2011). Antiproliferation and induction of apoptosis by *Moringa oleifera* leaf extract on human cancer cells. *Food Chem Toxicol*, **49**, 1270-5.
- Sultana S, Asif HM, Nazar HMI, et al (2014). Medicinal plants combating against cancer-a green anticancer approach. *Asian Pac J Cancer Prev*, **15**, 4385-94.
- Srisuwan S, Hamontri S, Kongsomboon K, et al (2014). See-and-treat approach to cervical intraepithelial lesions in HRH Princess Maha Chakri Sirindhorn Medical Center. *Asian Pac J Cancer Prev*, **15**, 3483-6.
- Tiloke C, Phulukdaree A, Chuturgoon AA (2013). The antiproliferative effect of *Moringa oleifera* crude aqueous leaf extract on cancerous human alveolar epithelial cells. *BMC Complement Altern Med*, **13**, 226.
- Vijayarathna S, Sreenivasan S (2012). Cytotoxicity of methanol extracts of *Elaeis guineensis* on MCF-7 and Vero cell lines. *Asian Pac J Trop Biomed*, **2**, 826-9.
- Vongsak B, Sithisarn P, Mangmool S, et al (2013a). Maximizing total phenolics, total flavonoids contents and antioxidant activity of *Moringa oleifera* leaf extract by the appropriate extraction method. *Indust Crops Products*, **44**, 566-71.
- Vongsak B, Sithisarn P, Gritsanapan W (2013b). Bioactive contents and free radical scavenging activity of *Moringa oleifera* leaf extract under different storage conditions. *Indust Crops Products*, **49**, 419-21.
- World Cancer Report (WCR) (2008). International agency for research on cancer. Retrieved 2011-02-26.